

Supplementary information

Structural and functional characterization of the Curli adaptor protein CsgF




Figure S1: Native (left) and denaturing/refolding (right) purification of His tagged CsgF analyzed by SDS-PAGE. M: protein marker; Full cell samples at 0h/2h time of protein expression; P_w: washed sample of insoluble cell debris after lysis; FT: flow through fraction of Ni-affinity chromatography; Wash: fractions of washing steps; RF: fraction of refolding step; Elution: fractions of elution steps. The eluted proteins were evaluated with MALDI mass spectrometry.




Figure S2: Far-UV spectra of CsgF purified under native and denaturing conditions. Spectra were recorded in 50 mM Phosphate Buffer at pH 7.

Table S1: Parameters of three dimensional NMR spectra, recorded at 600 MHz.

<i>Experiment</i>	<i>t1 (ms)</i>	<i>t2 (ms)</i>	<i>t3 (ms)</i>	<i>Mix (ms)</i>	<i>Scans</i>	<i>Ref.</i>
HNCO	17.6 (¹³ C)	17.9 (¹⁵ N)	121.7 (¹ H)	-	8	(Kay et al. 1990)
HN(CA)CO	22.0 (¹³ C)	20.9 (¹⁵ N)	121.7 (¹ H)	-	16	(Clubb et al. 1992)
CBCA(CO)NH	6.2 (¹³ C)	17.9 (¹⁵ N)	121.7 (¹ H)	-	12	(Grzesiek and Bax 1992)
HNCACB	7.6 (¹³ C)	17.9 (¹⁵ N)	121.7 (¹ H)	-	16	(Grzesiek and Bax 1992)
(H)CC(CO)NH	6.1 (¹³ C)	20.3 (¹⁵ N)	53.3 (¹ H)	20	32	(Montelione et al. 1992)
¹⁵ N edited TOCSY-HSQC	20.1 (¹ H)	20.3 (¹⁵ N)	56.8 (¹ H)	50	16	(Marion et al. 1989a)
¹³ C edited TOCSY-HSQC	20.9 (¹ H)	6.6 (¹³ C)	113.6 (¹ H)	60	8	(Bax et al. 1990)
¹³ C-HSQC-NOESY- ¹⁵ N-HSQC	6.4 (¹³ C)	20.3 (¹⁵ N)	60.9 (¹ H)	80	24	(Diercks et al. 1999)
¹⁵ N edited NOESY-HSQC	20.1 (¹ H)	28.7 (¹⁵ N)	56.8 (¹ H)	80	8	(Marion et al. 1989b)
¹³ C edited NOESY-HSQC (Ali)	20.9 (¹ H)	7.0 (¹³ C)	42.6 (¹ H)	80	8	(Marion et al. 1989b)
¹³ C edited NOESY-HSQC (Aro)	21.3 (¹ H)	6.0 (¹³ C)	85.2(¹ H)	80	4	(Marion et al. 1989b)




Figure S3: ^1H - ^{15}N HSQC Spectrum of 400 μM CsgF annotated with the resonance assignments (top), amino acid sequence of CsgF (bottom), unassigned residues are colored in red

Table S2: Statistics of NMR structure calculation and refinement.

<i>NMR constraints</i>	
Total NOE distance restraints	1027
Intraresidue, $i - j = 0$	368
Sequential, $i - j = 1$	386
Medium range, $1 < i - j < 5$	144
Long range, $i - j \geq 5$	129
Dihedral angle restraints	515
<i>Structural Statistics</i>	
Average violations per structure	
NOEs ≥ 0.1 Å	6 ± 2
Dihedrals $> 2^\circ$	19 ± 3
RMSD (Residues 68 – 127)	
Backbone (Å)	5.77 ± 1.75
Heavy (Å)	6.28 ± 1.77
<i>Ramachandran Analysis</i>	
Most favored region	65.2
Allowed region	28.6
Generously allowed	2.8
Disallowed	3.4
BMRB accession number	34052
PDB ID code	5M1U




Figure S4: ^1H - ^{15}N HSQC Spectra, Titration of paramagnetic substances.




Figure S5: Fitting of the Thioflavin T aggregation profiles using the unseeded (a) and the seeded (b) nucleation-elongation model of the Webtool Amylofit, applied to the first 3 hours of aggregation (a) or the full time frame (b)




Figure S6: Additional Thioflavin T amyloid aggregation assays (EB = Elution Buffer - CHES pH 10).




Figure S7: SDS-PAGE analysis of CsgB fibers aggregated with and without CsgF and washed with urea. M: marker; Bs: soluble CsgB serving as a control; F: CsgF; Bf: fibrillized CsgB. Washing fractions are numbered. Fibers were washed twice with 8 M urea/50 mM KPi pH 7.2 (steps 1 and 2) and two times in 125 mM Tris-HCl pH 7.5/0.8 M urea (steps 3 and 4). To dissolve CsgB fibers, the pellets were

resuspended in 6 M Guanidinium thiocyanate, 50 mM PBS pH 7.2 and precipitated with TCA (trichloroacetic acid).

CLUSTAL W (1.83) multiple sequence alignment

```

sp|CSGA_20-41    GVV-----PQYGGGGNHGGGGNNSG-----P
sp|CSGF_19-65    GTMTFQFRNPNFNGNPNGAFLNLSAQQAQNSYKDPSYNDDFGIETP
                  *.:          *::**.: *:.      **.*
CLUSTAL W (1.83) multiple sequence alignment

sp|CSGF_19-65    GTMTFQFRNPNFNGNPNGAFLNLSAQQAQNSYKDPSYNDDFGIETP---
sp|CSGB_21-41    AA-GYDLAN-----S-E-----YNFAVNELSKS
                  .:   ::: *                *   :                :*.:

```

Figure S8: Sequence Alignment of N-terminus of CsgF with CsgA or CsgB.

Supplementary References

- Bax A, Clore GM, Gronenborn AM (1990) ^1H - ^1H correlation via isotropic mixing of ^{13}C magnetization, a new three-dimensional approach for assigning ^1H and ^{13}C spectra of ^{13}C -enriched proteins. *J Magn Reson* 88:425–431. doi: 10.1016/0022-2364(90)90202-K
- Clubb RT, Thanabal V, Wagner G (1992) A constant-time three-dimensional triple-resonance pulse scheme to correlate intraresidue ^1HN , ^{15}N , and ^{13}C chemical shifts in $^{15}\text{N}/^{13}\text{C}$ -labelled proteins. *J Magn Reson* 97:213–217. doi: 10.1016/0022-2364(92)90252-3
- Diercks T, Coles M, Kessler H (1999) An efficient strategy for assignment of cross-peaks in 3D heteronuclear NOESY experiments. *J Biomol NMR* 15:177–180. doi: 10.1023/A:1008367912535
- Grzesiek S, Bax A (1992) Improved 3D triple-resonance NMR techniques applied to a 31 kDa protein. *J Magn Reson* 96:432–440. doi: 10.1016/0022-2364(92)90099-S
- Kay LE, Ikura M, Tschudin R, Bax A (1990) Three-dimensional triple-resonance NMR spectroscopy of isotopically enriched proteins. *J Magn Reson* 89:496–514. doi: 10.1016/0022-2364(90)90333-5
- Marion D, Driscoll PC, Kay LE, et al (1989a) Overcoming the overlap problem in the assignment of proton NMR spectra of larger proteins by use of three-dimensional heteronuclear proton-nitrogen-15 Hartmann-Hahn-multiple quantum coherence and nuclear Overhauser-multiple quantum coherence spectroscopy: application to interleukin 1. *Biochemistry* 28:6150–6156. doi: 10.1021/bi00441a004
- Marion D, Kay LE, Sparks SW, et al (1989b) Three-dimensional heteronuclear NMR of nitrogen-15 labeled proteins. *J Am Chem Soc* 111:1515–1517. doi: 10.1021/ja00186a066
- Montelione GT, Lyons BA, Emerson SD, Tashiro M (1992) An efficient triple resonance experiment using carbon-13 isotropic mixing for determining sequence-specific resonance assignments of isotopically-enriched proteins. *J Am Chem Soc* 114:10974–10975. doi: 10.1021/ja00053a051